

Consistency in Formulation of Spin Current and Torque Associated with a Variance of Angular Momentum

Yong Wang,^{1,2} Ke Xia,¹ Zhao-Bin Su,³ and Zhongshui Ma^{a2}

¹*Institute of Physics, Chinese Academy of Sciences, Beijing 100080, China*

²*School of Physics, Peking University, Beijing 100871, China*

³*Institute of Theoretical Physics, Chinese Academy of Sciences, Beijing 100080, China*

Stimulated generally by recent interest in the novel spin Hall effect, the nonrelativistic quantum mechanical conserved currents, taken into account of spin-orbit coupling, are rigorously formulated based on the symmetries of system and Noether's theorem. The quantum mechanical force on the spin as well as the torque associated with the variance of angular momentum are obtained. Consequently, the kinetic interpretation of the variances of spin and orbit angular momentum currents implies a torque on the "electric dipole" associated with the moving spin. The bearing of the force and the torque on the properties of spin current in a two-dimensional electron gas with the Rashba spin-orbit interaction is discussed.

PACS numbers: 72.10.Bg, 71.70.Ej, 72.25.-b

The theoretical prediction of spin Hall effect (SHE)[1, 2, 3, 4] has drawn considerable interest recently. The spin Hall current comes forward from the assembly effect that the carriers with opposite spins move in the opposite directions perpendicular to the electric field. Equivalently, a spin imbalance between its opposite flows is electrically generated via the spin-orbit (SO) coupling in the confined semiconductor. The manipulation of transverse force on the moving carriers with different spin orientations, i.e., pushes spin-up and -down carriers in the opposite directions, is understood as a dynamical issue of spintronics[5, 6, 7]. By using the optical methods, the experimental detections of spin accumulation in connection with SHE have recently been reported[8, 9]. The effect of impurity scattering on SHE has also been investigated[10, 11, 12].

However, a more fundamental question is emerged in the conventional understanding of SHE. The validity of the description for various physical properties of SHE depends on the correctness of definition for the spin current operator[13, 14, 15]. Since the translational and rotational motions are associated to the force and torque, without a rigorous definition of spin current, in some sense, the discussion about the convenient power source for the spin current is also somewhat arbitrary. Currently, the spin current operator is practically defined in analogy with that of charge current as a tensor operator of symmetric multiplying of Pauli matrix and velocity, $\mathbb{J}_\alpha^S = (\hbar/2)\{\sigma_\alpha, \mathbf{v}\}$ [2, 3, 4]. Nevertheless it does not satisfy a continuity equation[13, 14, 15], $\partial \rho_\alpha^S / \partial t + \nabla \cdot \mathbb{J}_\alpha^S \neq 0$, where $\rho_\alpha^S = (\hbar/2)\psi^\dagger \sigma_\alpha \psi$ is a spin density. Unlike the charge current, the conserved spin current can not be obtained within the context of Schrödinger equation with the addition of the SO interaction.

Actually, spin motion is intimately related to the orbital angular momentum of spin carriers. If we identify spin as the internal angular momentum of carriers, the orbit angular momentum can then be regarded as a dy-

namical property of carrier motion. The transmutation of spin orientation is compensated by that of orbit angular momentum during the carriers moving in the system. Although the spin current is not conserved, the total angular momentum consisted of both spin and orbit angular momenta should be conserved if there is no additional external force and torque.

In this paper, we inherit the Noether's idea[16], and provide a systematical formulation of the conservation laws for the translational and rotational motions of spin carriers based upon the generic symmetry principles. Considering a system with Dirac fermions coupled to an electromagnetic field, we formulated the practical forms of spin and orbit angular currents with the symmetries being maintained in the sense of the validity of Noether's theorem. The nonrelativistic approximation (NRA) of conserved currents is derived with the conservation laws being guaranteed. Under this formalism, the torque on the angular momentum and the force on spin carriers can be identified rigorously. As a result we gain an insight of torque on the transmutation of spin orientation and orbit angular momentum, and ascertain the role of SO coupling in the conservation laws.

Consider first the relativistically covariant Lagrangian of a closed system involved Dirac fields $\bar{\Psi}$, Ψ , and electromagnetic field A_μ . The corresponding Lagrangian is given by $\mathcal{L} = -(\hbar c/2)\bar{\Psi}\gamma_\mu \overleftrightarrow{\mathcal{D}}_\mu \Psi - mc^2\bar{\Psi}\Psi - (1/4)\mathcal{F}_{\mu\nu}\mathcal{F}_{\mu\nu}$ [16, 17], where the four-vector notations have been used ($\mu = 1, 2, 3, 4$): $x_\mu = (\mathbf{x}, ict)$, $A_\mu = (\mathbf{A}, i\varphi)$, and $\gamma_\mu = \left(\begin{pmatrix} 0 & -i\sigma_\alpha \\ i\sigma_\alpha & 0 \end{pmatrix}, \begin{pmatrix} I & 0 \\ 0 & -I \end{pmatrix} \right)$, σ_α ($\alpha = x, y, z$) are Pauli matrices; $f \overleftrightarrow{\mathcal{D}}_\mu g = f \mathcal{D}_\mu g - (\mathcal{D}_\mu^\dagger f)g$, $\mathcal{D}_\mu = \partial_\mu - i(e/\hbar c)A_\mu$ is a covariant derivation operator and $\mathcal{F}_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$ is the Maxwell field-strength tensor. From the Euler-Lagrangian partial differential equation, we can obtain the Dirac equation and the Maxwell's equation, $\gamma_\mu \mathcal{D}_\mu \Psi + (mc/\hbar)\Psi = 0$ and

$\partial_\nu \mathcal{F}_{\mu\nu} = ie\bar{\Psi}\gamma_\mu\Psi$, respectively.

According to Noether's theorem, any continuous one-parameter set of invariance of Lagrangian is associated with a local conserved current. For a closed system involved Dirac field and the electromagnetic field, the symmetries of system yield the conservation laws: (1) The U(1) symmetry leads to the charge conservation. Under the local U(1) gauge transformations, $\Psi \rightarrow \exp(-ie\Lambda/\hbar c)\Psi$, $\bar{\Psi} \rightarrow \bar{\Psi}\exp(ie\Lambda/\hbar c)$, and $A_\mu \rightarrow A_\mu - \partial_\mu\Lambda$, Noether's theorem gives rise to the continuity equation $\partial_\mu J_\mu = 0$ with the charge current density $J_\mu = iec\bar{\Psi}\gamma_\mu\Psi$; (2) Spatial translational isotropy begets the conservation of linear momentum while a time-translation symmetry gives rise to the energy conservation. The continuity equation $\partial_\mu \Theta_{\mu\nu} = 0$ is obtained from the invariance of Lagrangian under the time-space translation transformations $\delta x_\mu = \delta_{\mu\nu}a_\nu$, $\delta\Psi = 0$, $\delta\bar{\Psi} = 0$, and $\delta A_\mu = 0$. $\Theta_{\mu\nu} = (\hbar c/2)\bar{\Psi}\gamma_\mu\overleftrightarrow{D}_\nu\Psi + \mathcal{F}_{\mu\rho}\mathcal{F}_{\nu\rho} - \frac{1}{4}\mathcal{F}_{\mu\nu}\mathcal{F}_{\mu\nu} + \delta_{\mu\nu}$ is the energy-momentum current density tensor; (3) Rotational isotropy gives birth to the conservation of angular momentum. The Lagrangian is invariant under the transformations $x'_\mu = (\delta_{\mu\nu} + \epsilon_{\mu\nu})x_\nu$, $\Psi'(x') = [I - i(1/4)\epsilon_{\mu\nu}\sigma_{\mu\nu}]\Psi(x)$, $\bar{\Psi}'(x') = \bar{\Psi}(x)[I + i(1/4)\epsilon_{\mu\nu}\sigma_{\mu\nu}]$, and $A'_\mu(x') = (\delta_{\mu\nu} + \epsilon_{\mu\nu})A_\nu(x)$, with $\sigma_{\mu\nu} = i(1/2)[\gamma_\mu, \gamma_\nu]$. The total angular momentum current density tensor $J_{\lambda,\mu\nu} = \{x_\mu\Theta_{\lambda\nu} + \mathcal{F}_{\lambda\mu}A_\nu - (\mu \leftrightarrow \nu)\} - i(\hbar c/4)\bar{\Psi}\{\sigma_{\mu\nu}, \gamma_\lambda\}\Psi$ can be defined to satisfy the continuity equation $\partial_\lambda J_{\lambda,\mu\nu} = 0$.

The NRA of Dirac equation can be carried out by separating the Dirac wavefunction into large and small components[17], i.e., $\Psi = (\psi_A, \psi_B)^T \exp[-i(mc^2 + \epsilon)t/\hbar]$. Following the procedure of NRA given in Ref.[17], the normalized two-component wavefunction can be written in the form of $\psi = [1 - (\hbar^2/8m^2c^2)\sigma \cdot \mathbf{D}\sigma \cdot \mathbf{D}]\psi_A$ with a notation $\mathbf{D} = \nabla - i(e/\hbar c)\mathbf{A}$. The NRA Hamiltonian reads[17, 18] $H = -(\hbar^2/2m)\mathbf{D}^2 + e\varphi + i(e\hbar^2/4m^2c^2)\sigma \cdot (\mathbf{E} \times \mathbf{D}) - (e\hbar/2mc)\sigma \cdot \mathbf{B} - (e\hbar^2/8m^2c^2)\nabla \cdot \mathbf{E} - i(e\hbar^2/8m^2c^2)\sigma \cdot (\nabla \times \mathbf{E})$, where $\mathbf{E} = -\nabla\varphi$ and $\mathbf{B} = \nabla \times \mathbf{A}$ are the electric field and the magnetic field, respectively.

Alternative to obtain the continuity equation from Hamiltonian we engage the same procedure of NRA to the relativistic continuity equations. The corresponding NRA of currents can be obtained from the relativistic continuity equations by the power expansion in v^2/c^2 with the same procedure as obtaining the Hamiltonian with a SO coupling term. The symmetries of system are not broken by this power expansion. It is instructive to demonstrate it in the NRA of charge current $J_\mu = iec\bar{\Psi}\gamma_\mu\Psi$, which satisfies $\partial_\mu J_\mu = 0$ in the relativistic quantum mechanics. In the nonrelativistic limit to the order of v^2/c^2 , the continuity equation becomes $\dot{\rho}^C + \nabla \cdot \mathbf{J}^C = 0$. The charge density $\rho^C = e\psi^\dagger\psi + (\hbar^2e/8m^2c^2)\nabla^2(\psi^\dagger\psi) - i(\hbar^2e/8m^2c^2)\nabla \cdot (\psi^\dagger\sigma \times \overleftrightarrow{\mathbf{D}}\psi)$ and the current density $\mathbf{J}^C = -i(\hbar e/2m)\psi^\dagger\overleftrightarrow{\mathbf{D}}\psi +$

$(e^2\hbar/4m^2c^2)[\mathbf{E} + (2mc^2/e)\nabla] \times \psi^\dagger\sigma\psi$ have been expressed in terms of the basic local observables. Look at the charge density first. The second term concerns the variance of localized density of carriers. The last term is in connection with the accumulation of localized spin due to the spin flip. These two terms give a polarization vector

$$\mathbf{P} = -\frac{\hbar^2e}{8m^2c^2}[\nabla(\psi^\dagger\psi) - i\psi^\dagger\sigma \times \overleftrightarrow{\mathbf{D}}\psi]. \quad (1)$$

This shows that the nonuniform and the SO interaction in the system contribute an effective electric dipole. In the charge current the first two terms are recognized as identical to $e\rho^C\mathbf{v}$ with the Hamiltonian given above, where $\mathbf{v}$ is the velocity vector. The last term is a "spin magnetization current" $c\nabla \times \mathbf{m}$ arising from an "intrinsic magnetic moment", where $\mathbf{m} = (e\hbar/2mc)(\psi^\dagger\sigma\psi)$. The supplement of a magnetization term in the charge current has been established for a long time[19]. The Dirac equation implies the conservation law for the current $\partial_\mu J_\mu^C = 0$ as well as the decomposition $J_\mu^C = J_{\mu,G}^C + \partial_\nu m_{\mu\nu}$, where $J_{\mu,G}^C$ is Gordon current and $m_{\mu\nu} = i(e\hbar/4mc)\bar{\Psi}[\gamma_\mu, \gamma_\nu]\Psi$ is interpreted as the magnetic moment density due to the carrier spin[19].

The above analytic approach can be applied to the momentum current tensor $\Theta_{\mu\nu}$. From $\partial_\mu \Theta_{\mu\alpha} = 0$ we have the continuity equation $\dot{\rho}^M + \nabla \cdot \mathbb{J}^M = 0$, where the momentum density is given by

$$\rho^M = -i\frac{\hbar}{2}(1 + \frac{\hbar^2}{8m^2c^2}\nabla^2)\psi^\dagger\overleftrightarrow{\mathbf{D}}\psi + \rho_{SO}^M + \rho_G^M - \frac{\hbar^2e}{8m^2c^3}(\nabla\rho^C) \times \mathbf{B} + \frac{1}{c}\mathbf{E} \times \mathbf{B} \quad (2)$$

with $\rho_{SO}^M = -i(\hbar^2e/8m^2c^3)\psi^\dagger(\sigma\mathbf{B} \cdot \overleftrightarrow{\mathbf{D}} - \sigma \cdot \mathbf{B}\overleftrightarrow{\mathbf{D}})\psi$ and $\rho_G^M = (\hbar/16m^2c^2)\nabla \cdot (\psi^\dagger\sigma \times \overleftrightarrow{\mathbf{D}}\mathbf{D}\psi) - (\hbar^2e/8m^2c^3)\nabla \times (\psi^\dagger\psi\mathbf{B}) + (1/c)\nabla \cdot (\mathbf{E}\mathbf{A})$. $\mathbb{J}^M$ is the nonrelativistic limit of momentum current which takes a form as

$$\begin{aligned} \mathbb{J}^M = & -\frac{\hbar^2}{4m}\psi^\dagger\overleftrightarrow{\mathbf{D}}\mathbf{D}\psi - i\frac{\hbar^2e}{8m^2c^2}[\mathbf{E} \times (\psi^\dagger\sigma \overleftrightarrow{\mathbf{D}}\psi) \\ & + (\psi^\dagger\sigma \times \overleftrightarrow{\mathbf{D}}\psi)\mathbf{E} - \nabla(\psi^\dagger\psi)\mathbf{E} + (\nabla\mathbf{E})\psi^\dagger\psi] \\ & - (\mathbf{E}\mathbf{E} + \mathbf{B}\mathbf{B}) + \frac{1}{2}(E^2 + B^2 - \frac{\hbar e}{mc}\mathbf{B} \cdot \psi^\dagger\sigma\psi)\mathbb{I} \\ & + \frac{\hbar e}{2mc}\mathbf{B}\psi^\dagger\sigma\psi + \nabla \times (\mathbf{B}\mathbf{A} - i\frac{\hbar^2}{4m}\psi^\dagger\sigma \overleftrightarrow{\mathbf{D}}\psi) \end{aligned} \quad (3)$$

with the notation $\psi^\dagger\overleftrightarrow{D}_\beta D_\alpha\psi = \psi^\dagger D_\beta D_\alpha\psi + (D_\beta^*\psi^\dagger)\psi - (D_\alpha^*\psi^\dagger)(D_\beta\psi) - (D_\beta^*\psi^\dagger)(D_\alpha\psi)$.

The NRA of conservation of total angular momentum current tensor $J_{\lambda,\mu\nu}$ can be obtained in the same way. The local angular momentum density is found as

$$\rho^A = \tilde{\mathbf{x}} \times \rho^M + \rho^S + \frac{1}{c}\mathbf{E} \times \mathbf{A} + \frac{\hbar^2e}{4m^2c^3}\rho^C\mathbf{B}, \quad (4)$$

where $\tilde{\mathbf{x}} = \mathbf{x} - \mathbf{x}_0$ with a constant vector $\mathbf{x}_0$. The first term is the conventional angular momentum with the momentum density, while the last two terms in Eq. (4)

come from electromagnetic field. The second term is the intrinsic angular momentum density, associated with the local spin, given by $\rho^S = (\hbar/2)[1 - (\hbar^2/8m^2c^2)\nabla^2]\psi^\dagger\sigma\psi + \rho_{SO}^S + \rho_G^S$, where $\rho_{SO}^S = (\hbar^3/8m^2c^2)[2\psi^\dagger\sigma\mathbf{D}^2\psi + (\mathbf{D}^\dagger\psi^\dagger) \cdot \sigma(\mathbf{D}\psi) + (\mathbf{D}^\dagger\psi^\dagger)\sigma \cdot (\mathbf{D}\psi)]$ is the contribution of SO coupling induced by the transmutation and ρ_G^S relates to the surface terms $\nabla \cdot (\psi^\dagger\sigma\overleftrightarrow{\mathbf{D}}\psi)$ and $\nabla \times (\psi^\dagger\overleftrightarrow{\mathbf{D}}\psi)$. The local angular momentum density satisfies the continuity equation $\dot{\rho}^A + \nabla \cdot \mathbb{J}^A = 0$ with the total angular momentum current density tensor

$$\mathbb{J}^A = \tilde{\mathbf{x}} \times [\mathbb{J}^M + i\frac{\hbar^2}{4m}\nabla \times (\psi^\dagger\sigma\overleftrightarrow{\mathbf{D}}\psi)] - i\frac{\hbar^2}{4m}\psi^\dagger\overleftrightarrow{\mathbf{D}}\sigma\psi + i\frac{\hbar^2}{4m}\nabla \times [\psi^\dagger\sigma\overleftrightarrow{\mathbf{D}}\psi \times \tilde{\mathbf{x}}] + (\mathbf{B} \cdot \mathbf{A})\mathbb{I} - \mathbf{AB}, \quad (5)$$

where $\mathbb{I}$ is unit tensor. Notice that the survival of the term adhered the constant vector $\mathbf{x}_0$ in a conserved angular momentum of system is not a trivial matter. The continuity equation of angular momentum is satisfied with the symmetry of the invariance under spatial translation being guaranteed. Let us emphasize that the present form of total angular current is independent of the choice of coordinate origin in the system.

So far we have got the NRA expression of the conserved angular momentum current for a closed system including the fermion and the electromagnetic field. In general, the NRA conserved currents consist of three parts: orbital, spin, and electromagnetic. We next give a decision in favour of the angular momentum current of fermionic system by taking the electromagnetic field as an external field. Because the conservation laws are equations of motion for the observables, the force and the torque on the spin carriers, coupled to the external field, can be determined by the continuity equations. It is noted $\nabla \cdot [(\mathbf{E}\mathbf{E} + \mathbf{B}\mathbf{B}) - (1/2)(E^2 + B^2)\mathbb{I} + \nabla \times (\mathbf{B}\mathbf{A})] = \mathbf{E}\rho^C + (1/c)(\mathbf{J}^C \times \mathbf{B})$, which corresponds to Lorentz force, in the continuity equation of momentum flux when the Maxwell equation is applied to the time-independent situation. While the terms, in Eq.(3), $-(\hbar^2e/8m^2c^2)[i(\psi^\dagger\sigma \times \overleftrightarrow{\mathbf{D}}\psi)\mathbf{E} - \nabla(\psi^\dagger\psi)\mathbf{E} + (\nabla\mathbf{E})\psi^\dagger\psi] + (\hbar e/2mc)[\mathbf{B}\psi^\dagger\sigma\psi - (\mathbf{B} \cdot \psi^\dagger\sigma\psi)\mathbb{I}]$ can be regarded as the source of the force associated with the carrier's spin[5, 20]. The continuity equation of momentum current becomes $\dot{\rho}^{M'} + \nabla \cdot \mathbb{J}^{M'} = \mathbf{F}$, where the primes are meant to indicate the fermionic partition of density and current. The momentum density vector and the momentum current density tensor are given by $\rho^{M'} = -i(\hbar/2)\psi^\dagger\overleftrightarrow{\mathbf{D}}\psi$ and $\mathbb{J}^{M'} = -(\hbar^2/4m)\psi^\dagger\overleftrightarrow{\mathbf{D}}\overleftrightarrow{\mathbf{D}}\psi - i(\hbar^2e/8m^2c^2)\mathbf{E} \times (\psi^\dagger\sigma\overleftrightarrow{\mathbf{D}}\psi) - i(\hbar^2/4m)\nabla \times (\psi^\dagger\sigma\overleftrightarrow{\mathbf{D}}\psi)$, respectively. The total force is found

$$\mathbf{F} = e(\mathbf{E} + \frac{\hbar^2}{8m^2c^2}\nabla^2\mathbf{E})\psi^\dagger\psi + [\mathbf{J}^C - c\nabla \times \mathbf{m}] \times \mathbf{B} + \frac{\hbar e}{2mc}(\nabla\mathbf{B}) \cdot \rho^S + i\frac{\hbar^2e}{8m^2c^2}(\nabla\mathbf{E}) \cdot \psi^\dagger\sigma \times \overleftrightarrow{\mathbf{D}}\psi. \quad (6)$$

The first three terms are the Lorentz force and the "Stern-Gerlach force" which relates to a magnetic field gradient $\nabla\mathbf{B}$. The forth term is new which has not been discussed previously. It relates to the fluctuation of localized spin current and produces a torque on the angular momentum. The continuity equation of angular momentum has the general form $\dot{\rho}^{A'} + \nabla \cdot \mathbb{J}^{A'} = \mathbf{M}$, where $\mathbb{J}^{A'}$ is the total angular momentum current of electron which can be written in the form of a sum of three terms $\mathbb{J}^{A'} = \mathbb{J}^O + \mathbb{J}^S + i(\hbar^2/4m)\nabla \times (\psi^\dagger\sigma\overleftrightarrow{\mathbf{D}}\psi \times \tilde{\mathbf{x}})$. The orbit angular momentum current is given by

$$\mathbb{J}^O = \tilde{\mathbf{x}} \times [\mathbb{J}^{M'} - i\frac{\hbar^2}{4m}\nabla \times (\psi^\dagger\sigma\overleftrightarrow{\mathbf{D}}\psi)] \quad (7)$$

and the spin current is defined by

$$J_{\beta\alpha}^S = -i\frac{\hbar^2}{4m}\psi^\dagger\sigma_\alpha\overleftrightarrow{D}_\beta\psi - \epsilon_{\beta\alpha\gamma}\frac{\hbar^2e}{8m^2c^2}E_\gamma\psi^\dagger\psi. \quad (8)$$

The term on the right hand side of continuity equation is the torque $\mathbf{M}$ which relates to the total angular momentum, not spin partition itself. We find

$$\mathbf{M} = \tilde{\mathbf{x}} \times \mathbf{F} + \frac{e}{mc}\rho^S \times \mathbf{B} + i\frac{\hbar^2e}{8m^2c^2}(\psi^\dagger\sigma \times \overleftrightarrow{\mathbf{D}}\psi) \times \mathbf{E}. \quad (9)$$

There are several origins in the torque $\mathbf{M}$. Besides the conventional torque caused by the force $\mathbf{F}$, the second term is the torque in the fundamental equation of spin resonance theory and the third term is a torque induced by the spin current. Because we are ultimately interested in relating the induced torque to the effect of SO coupling, a fluctuation in the spin current $\Delta\mathbf{J}^S = i(\hbar^2/4m)\psi^\dagger(\sigma \times \overleftrightarrow{\mathbf{D}})\psi$ is introduced or, equivalently

$$\Delta J_\alpha^S = \epsilon_{\alpha\beta\gamma}J_{\beta\gamma}^S + \frac{\hbar^2e}{4m^2c^2}E_\alpha\psi^\dagger\psi. \quad (10)$$

By using Eq. (8) the relation between ΔJ_α^S and the spin current can be put in the form $\epsilon_{\alpha\beta\gamma}\Delta J_\gamma^S = J_{\alpha\beta}^S - J_{\beta\alpha}^S$. This vector has its own significance[21]. It contributes a force if there is a spatial change of electric field and a torque on the spin when the carriers moves in an electric field. From Eqs. (6) and (9) an force $\Delta\mathbf{F}$ and a torque $\Delta\mathbf{M}$ induced by the SO coupling may be obtained, which can further be separated into the contributions from the accumulation of spin and the fluctuation of spin current, in a compact form

$$\begin{pmatrix} \Delta\mathbf{F} \\ \Delta\mathbf{M} \end{pmatrix} = \frac{e}{2mc^2} \begin{pmatrix} 2c(\nabla\mathbf{B}) \cdot & (\nabla\mathbf{E}) \cdot \\ -2c\mathbf{B} \times & -\mathbf{E} \times \end{pmatrix} \begin{pmatrix} \rho^S \\ \Delta\mathbf{J}^S \end{pmatrix}. \quad (11)$$

It is noted that $\Delta\mathbf{J}^S$ is just the "electric dipole" $\langle \hat{\mathbf{I}} \rangle = -\langle (\hat{\mu}/2c) \times (-i\hbar\hat{\mathbf{D}}/m) \rangle$ with the intrinsic magnetic dipole moment $\hat{\mu} = (\hbar e/2mc)\sigma$. Physically, it is thinking to be a spin vector rotates around an axis with a velocity

$-i\hbar(\mathbf{D}/m)$ to give rise a magnetic current circuit, equivalently, an "electric dipole". The torque on the "electric dipole" is $\langle \hat{\mathbf{I}} \rangle \times \mathbf{E}$ as expected.

Next, we apply the formulism to a two-dimensional electron gas (2DEG) with the Rashba SO interaction. This can be achieved by taking $\mathbf{E} = (0, 0, E_z)$ and $\mathbf{A} = 0$ in our derivations. In the follow discussion the indices k and l are specialized to 2D indices while α still is indicated in 3D space. The Rashba Hamiltonian is then given by $H_R = (1/2m)\mathbf{p}^2 - (\lambda/\hbar)(p_x\sigma_y - p_y\sigma_x)$ with $\lambda = (\hbar^2 e / 4m^2 c^2) E_z$. The momentum current in Eq. (3) reduces to $J_{kl}^{\mathcal{M}'} = -(\hbar^2/4m)\psi^\dagger \overleftrightarrow{\nabla}_k \overleftrightarrow{\nabla}_l \psi + i\epsilon_{kjlz}(\lambda/2)[\psi^\dagger \sigma_j \overleftrightarrow{\nabla}_l \psi - (\hbar^2/2m\lambda)\partial_j(\psi^\dagger \sigma_z \overleftrightarrow{\nabla}_l \psi)]$. It is easy to prove that $\nabla_k J_{kl}^{\mathcal{M}'} = 0$ and $F_\beta = 0$ for the stationary states. The total angular momentum current is given by

$$J_{k\alpha}^{A'} = J_{k\alpha}^{\mathcal{O}} + J_{k\alpha}^{\mathcal{S}} + i\frac{\hbar^2}{4m}\nabla \times [(\psi^\dagger \sigma \overleftrightarrow{\mathbf{D}} \psi) \times \tilde{\mathbf{x}}]_{k\alpha}. \quad (12)$$

It is shown that the orbit part of angular momentum current density, $J_{k\alpha}^{\mathcal{O}} = -(\hbar^2/4m)\epsilon_{\alpha jl}x_j(\psi^\dagger \overleftrightarrow{\nabla}_l \overleftrightarrow{\nabla}_k \psi) - i(\lambda/2)\delta_{\alpha z}\tilde{\mathbf{x}} \cdot (\psi^\dagger \sigma \overleftrightarrow{\nabla}_k \psi)$, is nonvanishing only for $\alpha = z$. The corresponding spin current is given by $J_{k\alpha}^{\mathcal{S}} = -i(\hbar^2/4m)\psi^\dagger \sigma_\alpha \overleftrightarrow{\nabla}_k \psi - \epsilon_{k\alpha z}(\lambda/2)(\psi^\dagger \psi)$. It is shown that the divergence of spin current, in general, satisfies the equation $\dot{\rho}^{\mathcal{S}} + \nabla \cdot \mathbb{J}^{\mathcal{S}} = \mathbf{T}$ with the torque $T_{x(y)} = i(\lambda/2)(\psi^\dagger \sigma_z \overleftrightarrow{\nabla}_{x(y)} \psi)$ and $T_z = -i(\lambda/2)(\psi^\dagger \sigma \cdot \overleftrightarrow{\nabla} \psi)$. A straightforward calculation shows $\nabla_k J_{kz}^{\mathcal{O}} = -T_z$. This leads to $\nabla_k J_{kz}^{A'} = 0$, while $\nabla_k J_{kl}^{A'} = T_l$, i.e. the component of total angular momentum perpendicular to the x-y plane is conserved, which shows clearly how the transmutation between the spin and the orbit angular momenta occurs in the system with a SO coupling, while the in-plan components of spin current are not conserved by requiring a torque on it. It is worth to emphasize that $\mathbf{F}$ is nonvanishing in the presence of an in-plan electric field. The force gives rise to an extra conventional torque. In general, it can be shown a relation between the torques $\mathbf{T}$ and $\Delta\mathbf{M}$, acting on the spin itself and on the total angular momentum, respectively,

$$\Delta\mathbf{M} = \mathbf{T} + i\frac{\hbar^2 e}{8m^2 c^2}\psi^\dagger(\sigma \times \mathbf{E}) \times \overleftrightarrow{\mathbf{D}} \psi. \quad (13)$$

We note that there is an essential difference between the force $\mathbf{F}$ defined from the continuity equation of momentum current and the transverse force $\mathbf{f}$ on the spin for driving spin current. The latter is the covariant force on a spin defined from the equation of motion

$$m\dot{\mathbf{v}} = -m\nabla \cdot \mathbb{J}^v + \mathbf{f}, \quad (14)$$

where $\mathbf{v}$ is the velocity of a wave packet of spin-polarized electron. In a Rashba 2DEG it is found $v_j = -(\hbar^2/2m)\psi^\dagger \overleftrightarrow{D}_j \psi - (2\lambda/\hbar^2)\epsilon_{zjk}\rho_k^{\mathcal{S}} + (1/m)\epsilon_{jkl}\nabla_k \rho_l^{\mathcal{S}}$. $\mathbb{J}^v$

is the corresponding tensor of velocity flux given by $J_{jk}^v = (1/m)J_{jk}^{\mathcal{M}'} + (2\lambda/\hbar^2)\epsilon_{zlk}J_{jl}^{\mathcal{S}}$. The second term in the velocity flux can be understood as a drift velocity because of bringing the electric field to bear on the moving spin. The covariant force on a spin is found as

$$f_j = -\frac{2m\lambda}{\hbar^2}\epsilon_{zjk}T_k + \epsilon_{jk\alpha}\nabla_k T_\alpha. \quad (15)$$

Replacing $T_k = -(2m\lambda/\hbar^2)J_{kz}^{\mathcal{S}}$ into the second term of f_j , we find $f_j = (4m^2\lambda^2/\hbar^4)\epsilon_{zjk}J_{kz}^{\mathcal{S}}$. Because of the opposite signs for the contrary components of spin-polarized spin current, the different spins experience the transverse forces in the opposite directions[6, 7].

The main results of this paper may now be summarized as follows. The conservation law of NRA conserved angular momentum based on the Noether's theorem has been formulated systematically and rigorously. It is identified with that the torques are generated on a moving spin through the "electric dipole" when the electromagnetic field is regarded as an external field. As an example of the application, we consider a Rashba 2DEG and show that a torque will develop on the spin. The transmutation between the spin and the orbit angular momenta has been demonstrated explicitly. Furthermore, the covariant force on the spin in the presence of SO coupling is clarified.

This work is supported by NNSFC grants 10274069 and 10474002.

^{a)} Electronic mail: mazs@phy.pku.edu.cn

-
- [1] J.E. Hirsch, Phys. Rev. Lett. **83**, 1834(1999)
 - [2] S. Murakami, N. Nagaosa, and S.C. Zhang, Science **301**, 1348(2003)
 - [3] J. Sinova, D. Culcer, Q. Niu, N.A. Sinitsyn, T. Jungwirth, and A.H. MacDonald, Phys. Rev. Lett. **92**, 126603(2004)
 - [4] S.Q. Shen, M. Ma, X.C. Xie, and F.C. Zhang, Phys. Rev. Lett. **92**, 256603(2004)
 - [5] C.M. Ryu, Phys. Rev. Lett. **76**, 968(1996)
 - [6] B.K. Nikolić, L.P.Zárbo, and S. Welack, cond-mat/0503415
 - [7] S.Q. Shen, cond-mat/0506676
 - [8] Y.K. Kato, R.C. Myers, A.C. Gossard, and D.D. Awschalom, Science **306**, 1950(2004)
 - [9] J. Wunderlich, B. Kaestner, J. Sinova, and T. Jungwirth, Phys. Rev. Lett. **94**, 047204(2005)
 - [10] J.I. Inoue, G.E. Bauer, and L.W. Molenkamp, Phys. Rev. B **70**, 041303(R)(2004)
 - [11] E.G. Mishchenko, A.V. Shytov, and B.I. Halperin, Phys. Rev. Lett. **93**, 226602(2004)
 - [12] L. Sheng, D.N. Sheng, and C.S. Ting, Phys. Rev. Lett. **94**, 016602(2005)
 - [13] Q.F. Sun and X.C. Xie, cond-mat/0502317
 - [14] P. Zhang, J.R. Shi, D. Xiao, and Q. Niu, cond-mat/0503505
 - [15] J. Wang, B.G. Wang, W. Ren, and H. Guo, cond-mat/0507159

- [16] C. Itzykson and J.-B. Zuber, Quantum Field Theory, McGraw-Hill Inc. (1980)
- [17] J.J. Sakurai, Advanced Quantum Mechanics, Addison-Wesley Publishing Company (1967)
- [18] R. Winkler, Physica **E22**, 450(2004)
- [19] R. Gurtler and D. Hestenes, J. Math. Phys. **16**, 573(1975)
- [20] S.R. Barone, Phys. Rev. D **8**, 3492(1973)
- [21] E.I. Rashba, Phys. Rev. B **68**, 241315(R)(2003)